



W. Garrett Owen¹
wgowen@msu.edu

Ornamental Sweetpotato Intumescence: *A Physiological Disorder*

Sporadic green bumps and clusters of translucent or white wart-like lesions were recently observed among veins of dark-leaf ornamental sweetpotato vines. These abnormalities growing on the leaf surface are a result of a physiological disorder, intumescence.

During a recent greenhouse visit, I inspected a crop of dark purple ornamental sweetpotato (*Ipomoea batatas*) plants. The matured leaves were exhibiting abnormal growth on the upper surface (Fig. 1). The growth ranged from sporadic green bumps to translucent or white lesions to black out-growths (Fig. 2).

At closer examination, the small green bumps were rising between the veins of the leaf and along the mid-rib. Though the green bumps were intermittent, I noticed a pattern among the leaves. I found some leaves to only exhibit small green bumps, while others, the green bumps were starting to enlarge and extrude out from the leaf surface, turning translucent (Fig. 3). The enlarged translucent lesions appeared white due to the intensity of out-growth (Fig. 4) that were present along or on the mid-rib and

2017 Sponsors



Figure 1. Mature leaves of sweetpotato vines exhibiting abnormal, translucent out-growths on the upper leaf surface.

e-GRO Alert

www.e-gro.org

CONTRIBUTORS

Dr. Nora Catlin
 Floriculture Specialist
 Cornell Cooperative Extension - Suffolk County
 nora.catlin@cornell.edu

Dr. Chris Currey
 Assistant Professor of Floriculture
 Iowa State University
 ccurrey@iastate.edu

Dr. Ryan Dickson
 Ext. Specialist for Greenhouse Management & Technologies
 University of New Hampshire
 ryan.dickson@unh.edu

Thomas Ford
 Commercial Horticulture Educator
 Penn State Extension
 tgf2@psu.edu

Dan Gilrein
 Entomology Specialist
 Cornell Cooperative Extension - Suffolk County
 dog1@cornell.edu

Dr. Joyce Latimer
 Floriculture Extension & Research
 Virginia Tech
 jlatime@vt.edu

Dr. Roberto Lopez
 Floriculture Extension & Research
 Michigan State University
 rglopez@msu.edu

Dr. Neil Mattson
 Greenhouse Research & Extension
 Cornell University
 neil.mattson@cornell.edu

Dr. Garrett Owen
 Floriculture Outreach Specialist - Michigan State Univ.
 wgowen@msu.edu

Dr. Rosa E. Raudales
 Greenhouse Extension Specialist
 University of Connecticut
 rosa.raudales@uconn.edu

Dr. Beth Scheckelhoff
 Ext. Educator – Greenhouse Systems
 The Ohio State University
 scheckelhoff.11@osu.edu

Lee Stivers
 Extension Educator – Horticulture
 Penn State Extension, Washington County
 ljs32@psu.edu

Dr. Paul Thomas
 Floriculture Extension & Research
 University of Georgia
 pathomas@uga.edu

Dr. Ariana Torres-Bravo
 Horticulture/ Ag. Econ., Purdue University
 torres2@purdue.edu

Dr. Brian Whipker
 Floriculture Extension & Research - NC State Univ.
 bwhipker@ncsu.edu

Heidi Wollaeger
 Floriculture Outreach Specialist - Michigan State Univ.
 wollaeger@anr.msu.edu

Copyright © 2017


Where trade names, proprietary products, or specific equipment are listed, no discrimination is intended and no endorsement, guarantee or warranty is implied by the authors, universities or

most of the growth occurring at the leaf base (Fig. 5).

In some instances, I found the out-growth to be black (Fig. 6). While the majority of the abnormal out-growth occurred on the upper leaf surface, I did find some growth arising from the leaf petiole. The abnormal growth is a physiological disorder termed, intumescence, however other common and interchangeable names include: excrescences, neoplasms, galls, genetic tumors, lesions, enations, and oedemata.

Causative Factors

Intumescence development on leaves and petioles can have an impact on the aesthetic value of ornamental crops. You may be wondering, “What causes this physiological disorder?” The causative factor related to intumescence development is rather vague. Many have proposed intumescence development to be a result of air contamination, carbohydrate balance, chemical application, excess water, genetics, hormones and hormone concentration, humidity, light quality and quantity (intensity), and temperature. To date, there is no cure for intumescence development on ornamental sweetpotato. The best method to avoid intumescence



© W. Garrett Owen

Figure 2. Abnormal growth on the upper leaf ranged from sporadic green bumps to translucent or white lesions to black out-growths.

development on ornamental sweetpotato crops would be to select cultivars that are less susceptible. Trials conducted at Kansas State University determined which ornamental sweetpotato cultivars were the least and most susceptible to intumescence development (Table 1). For more information, watch the two e-GRO webinars: Blisters, Bumps and Lesions: The Physiological Disorders of Intumescence and Edema ([Link to Part 1](#)) and ([Link to Part 2](#)).

Literature Cited

Craver, J. K., C.T. Miller, M.G. Cruz, K.A. Williams. 2014. Intumescences: Further Investigations into an Elusive Physiological Disorder. Greenhouse Production News. 24(9):32-40.



Figure 3. Small green bumps protrude out from the leaf surface, turning translucent or white along the mid-rib.

Cooperating Universities

UConn



Cornell University



The University of Georgia

IOWA STATE UNIVERSITY

MICHIGAN STATE UNIVERSITY

NC STATE UNIVERSITY



THE OHIO STATE UNIVERSITY

PENNSTATE



Cooperative Extension
College of Agricultural Sciences

PURDUE UNIVERSITY



University of New Hampshire

Cooperative Extension



VirginiaTech
Invent the Future®

MAUMEE VALLEY GROWERS
Choose the Very Best.



In cooperation with our local and state greenhouse organizations



Indiana FLOWER GROWERS Association



Michigan Floriculture Growers Council



CONNECTICUT GREENHOUSE GROWERS ASSOCIATION



© W. Garrett Owen

Figure 4. Microscope view of enlarged translucent lesions present along the mid-rib of an ornamental sweetpotato leaf.



© W. Garrett Owen

Figure 5. Enlarged translucent lesions appear white due to the intensity of out-growth that are present along or on the mid-rib and with most of the growth occurring at the leaf base.



Figure 6. Translucent out-growth turn black and become dry.

Table 1. List of ornamental sweetpotato (*Ipomoea batatas*) cultivars screened in a trial conducted at Kansas State University (Craver et al., 2014).

Symptomatic Cultivars (cultivars having less than 5% leaves affected by intumescence)			
	Leaf Color	Plant Habit	Leaf Shape
'Bright Ideas Black'	deep-purple	trailing	palmate, longer, thin lobes
'Bright Ideas Lime'	yellow-green	compact	palmate, shorter, thick lobes
'Bright Ideas Rusty Red'	bronze-red, green	compact	shield-shaped
'Desana Compact Red'	bronze-red, green	trailing	palmate, shorter, thick lobes
'Sidekick Black'	deep-purple	compact	palmate, longer, thin lobes
'Sweet Caroline Green Yellow'	green with yellow/white streaking	compact	palmate, longer, thin lobes
'Sweet Caroline Sweetheart Purple'	deep-purple	trailing	heart-shaped
'Sweet Georgia Bronze'	bronze-red, green	compact	shield-shaped
'Sweet Georgia Heart Light Green'	yellow-green	compact	heart-shaped
Highly Symptomatic Cultivars (cultivars having more than 20% leaves affected by intumescence)			
'Black Heart'	deep-purple	trailing	heart-shaped
'Blackie'	deep-purple	trailing	palmate, longer, thin lobes
'Desana Bronze'	red-bronze	trailing	heart-shaped
'South of the Border Chipotle'	deep-purple with green splotches	trailing	heart-shaped
'Sweet Caroline Bronze'	bronze-red	trailing	palmate, shorter, thick lobes
'Sweet Caroline Sweetheart Light Green'	yellow-green	compact	heart-shaped
'Sweet Caroline Sweetheart Red'	bronze-red, green	trailing	heart-shaped
'Tricolor'	pink-white-green	compact	shield-shaped
Non-Symptomatic Cultivars			
'Desana Lime'	yellow-green	trailing	heart-shaped
'FloraMia Nero'	deep-purple	compact	palmate, shorter, thick lobes
'Illusion Emerald Lace'	yellow-green	trailing	palmate, longer, very thin lobes
'Illusion Garnet Lace'	bronze-red	compact	palmate, longer, very thin lobes
'Illusion Midnight Lace'	deep-purple	trailing	palmate, longer, very thin lobes
'Margarita'	yellow-green	trailing	heart-shaped
'Sidekick Black Heart'	deep-purple	compact	heart-shaped
'Sidekick Lime'	yellow-green	compact	heart-shaped
'South of the Border Chihuahua'	yellow-green	compact	palmate, shorter, thick lobes
'South of the Border Guacamole'	green-bronze	trailing	palmate, shorter, thick lobes
'Sweet Caroline Bewitched'	deep-purple	compact	shield-shaped
'Sweet Caroline Light Green'	yellow-green	trailing	palmate, shorter, thick lobes
'Sweet Caroline Raven'	deep-purple	compact	palmate, shorter, thick lobes
'Sweet Caroline Red'	bronze-red	trailing	palmate, shorter, thick lobes
'Sweet Georgia Bullfrog'	deep-purple with green splotches	compact	palmate, longer, thin lobes
'Sweet Georgia Heart Deep Purple'	deep-purple	compact	palmate, shorter, thick lobes
'Sweet Georgia Heart Purple'	deep-purple	trailing	heart-shaped
'Sweet Georgia Heart Red'	bronze-red, green	compact	heart-shaped
'Sweet Georgia Light Green'	yellow-green	compact	palmate, shorter, thick lobes

(Craver et al., 2014)