



Hunter Landis<sup>1</sup>

Brian E. Whipker<sup>2</sup>  
bwhipker@ncsu.edu

## Hydrangea Nutrition: Upper Leaf Interveinal Chlorosis (Iron Deficiency)

*Interveinal chlorosis (yellowing) of the upper foliage is an indicator that iron (Fe) is limited in availability. Many causal factors can influence iron availability and are discussed in this Alert.*

Interveinal chlorosis is readily identified as a problem associated with inadequate levels of iron (Fe) (Fig. 1). Many factors affect Fe uptake. In greenhouse production, elevated substrate pH, which makes Fe unavailable to the plant, is usually the primary suspect (Fig. 2).

Also keep in mind a number of other factors can also negatively impact Fe uptake by plants. Hydrangeas (*Hydrangea macrophylla*) are often grown with cooler temperatures. As a standard rule, substrate temperatures below 55F (13C) limits Fe uptake by the plant.

In addition, at these lower temperatures, water use by the plant can be less, thus making the plants more susceptible to overwatering. Iron (and phosphorus) uptake is a challenge for hydrangeas, like most plants, with excessively wet conditions. Prolonged substrate wetness inhibits root development, which can

<sup>1</sup> Hunter Landis is an Agronomist at the North Carolina Department of Agriculture – Plant, Waste, Solution and Media section. His responsibilities include providing interpretation guidelines for tissue analysis results. <sup>2</sup> Brian Whipker is a professor of floriculture at North Carolina State University.

2017 Sponsors



Figure 1. Interveinal chlorosis of the upper leaves most times indicate that iron is limited.

**e-GRO Alert**

*www.e-gro.org*

**CONTRIBUTORS**

**Dr. Nora Catlin**  
 Floriculture Specialist  
 Cornell Cooperative Extension - Suffolk County  
 nora.catlin@cornell.edu

**Dr. Chris Currey**  
 Assistant Professor of Floriculture  
 Iowa State University  
 ccurrey@iastate.edu

**Dr. Ryan Dickson**  
 Ext. Specialist for Greenhouse Management & Technologies  
 University of New Hampshire  
 ryan.dickson@unh.edu

**Thomas Ford**  
 Commercial Horticulture Educator  
 Penn State Extension  
 tgf2@psu.edu

**Dan Gilrein**  
 Entomology Specialist  
 Cornell Cooperative Extension - Suffolk County  
 dog1@cornell.edu

**Dr. Joyce Latimer**  
 Floriculture Extension & Research  
 Virginia Tech  
 jlatime@vt.edu

**Dr. Roberto Lopez**  
 Floriculture Extension & Research  
 Michigan State University  
 rglopez@msu.edu

**Dr. Neil Mattson**  
 Greenhouse Research & Extension  
 Cornell University  
 neil.mattson@cornell.edu

**Dr. Garrett Owen**  
 Floriculture Outreach Specialist - Michigan State Univ.  
 wgowen@msu.edu

**Dr. Rosa E. Raudales**  
 Greenhouse Extension Specialist  
 University of Connecticut  
 rosa.raudales@uconn.edu

**Dr. Beth Scheckelhoff**  
 Ext. Educator – Greenhouse Systems  
 The Ohio State University  
 scheckelhoff.11@osu.edu

**Lee Stivers**  
 Extension Educator – Horticulture  
 Penn State Extension, Washington County  
 ljs32@psu.edu

**Dr. Paul Thomas**  
 Floriculture Extension & Research  
 University of Georgia  
 pathomas@uga.edu

**Dr. Ariana Torres-Bravo**  
 Horticulture/ Ag. Econ., Purdue University  
 torres2@purdue.edu

**Dr. Brian Whipker**  
 Floriculture Extension & Research - NC State Univ.  
 bwhipker@ncsu.edu

**Heidi Wollaeger**  
 Floriculture Outreach Specialist - Michigan State Univ.  
 wollaeger@anr.msu.edu


Copyright © 2017

*Where trade names, proprietary products, or specific equipment are listed, no discrimination is intended and no endorsement, guarantee or warranty is implied by the authors, universities or*

exacerbate the situation already occurring in the plants.

The impact of these cultural factors of cool temperatures and saturated substrates can be managed by careful water management. In practice it may be difficult due to plant-to-plant variation. One will need to irrigate when the majority of the crop requires water. If possible, the plants that dry out slower may need to be grouped and irrigated at a less frequent interval.

From a plant diagnostic perspective, symptoms are often visible on a larger percentage of the crop when Fe is limited due to low temperatures or overall wet substrate conditions (Fig. 3). If a Fe deficiency is suspected, then submit a substrate and tissue sample to a commercial lab for confirmation.



© Brian Whipker

Figure 2. The interaction of substrate pH on nutrient availability. At higher pH levels greater than 6.5, iron is tied up.


Root rot is another factor that needs to be considered if upper leaf interveinal chlorosis is occurring. Root rot problems usually occur on scattered, individual plants and the new growth is usually stunted. So the scattered pattern of stunted plants will aid in your diagnosis of a root rot problem. In addition, when leaf interveinal chlorosis occurs, always inspect the roots. Black or brown roots are a strong indicator of rot. Submit a sample to a disease diagnostic clinic to confirm your diagnosis.

In this situation we analyzed the interveinal chlorotic symptomatic leaves and a sample from dark green leaves as a “good” comparison. The tissue analysis results confirmed that Fe was limited (Fig. 4). The good comparison plants had 121 ppm Fe, while the leaves with initial to moderate interveinal chlorosis symptoms contained 78.4 ppm Fe. Plants with advanced symptoms of bright yellow leaves contained 37.7 ppm Fe in the tissue. The recommended range for Fe varies by source. Bryson and Mills (2014) list a range between 50 and 300 ppm Fe, while Dole and Wilkins (2005) have a narrower 85 to 115 ppm Fe range. The plants exhibiting initial to moderate upper leaf interveinal chlorosis were within the acceptable range suggested by Bryson and Mills, but below the recommended range of Dole and Wilkins. The leaves with more advanced symptoms were below the recommended ranges of both references.

In this situation of observing interveinal chlorotic leaves scattered throughout the crop, root rot was suspected. The roots were then inspected and were white and healthy, but limited root growth was noticed. The interveinal chlorosis was likely confounded due to the previous week of cloudy weather, which limited the frequency when fertilizer was applied, and resulted in prolonged substrate wetness.

Cooperating Universities

UConn



Cornell University



The University of Georgia

IOWA STATE UNIVERSITY

MICHIGAN STATE UNIVERSITY

NC STATE UNIVERSITY



THE OHIO STATE UNIVERSITY

PENNSTATE



Cooperative Extension  
College of Agricultural Sciences

PURDUE UNIVERSITY



University of New Hampshire  
Cooperative Extension



VirginiaTech  
Invent the Future®

MAUMEE VALLEY GROWERS  
Choose the Very Best.



In cooperation with our local and state greenhouse organizations



Indiana FLOWER GROWERS Association



Michigan Floriculture Growers Council



CONNECTICUT GREENHOUSE GROWERS ASSOCIATION



Figure 3. Interveinal chlorosis of the upper leaves of only a few plants suggest either a cultural problem or root rot is occurring. Inspect the roots for poor growth or root rot to help diagnose the situation.

We took one of the plants back to our greenhouse to grow it longer and observe the symptoms. By customizing the irrigation frequency based on the need of this individual plant, the plant was able to recover on its own without applying any acidification corrective procedures. The leaves turned green within 2 weeks.

This example illustrates the interaction between slow root growth, environmental conditions, and excessive irrigations. With plant growth, it is impossible to decouple them, thus in order to avoid this situation, all these factors must be addressed when taking corrective steps.




Figure 4. A comparison of tissue iron (Fe) values from a lab analysis. Low Fe levels were confirmed by the test.

**Literature Cited:**

Bryson, G.M. and H.A. Mills (editors). 2014. Plant Analysis Handbook IV, Micro-Macro Publishing, Athens Georgia. p. 600.

Dole, J.M. and H.F. Wilkins. 2005. Floriculture Principles and Species, 2nd Ed. Pearson-Prentice Hall, Upper Saddle River, NJ. p. 1023.

© Brian Whipker